



## Anthropogenic debris on beaches in the SE Pacific (Chile): Results from a national survey supported by volunteers

Macarena Bravo<sup>a</sup>, M<sup>a</sup> de los Ángeles Gallardo<sup>a</sup>, Guillermo Luna-Jorquera<sup>a,b</sup>, Paloma Núñez<sup>a</sup>,  
Nelson Vásquez<sup>a</sup>, Martin Thiel<sup>a,b,\*</sup>

<sup>a</sup> Universidad Católica del Norte, Larrondo 1281, Coquimbo, Chile

<sup>b</sup> Centro de Estudios Avanzados en Zonas Áridas (CEAZA), Coquimbo, Chile

### ARTICLE INFO

#### Keywords:

Beach litter  
Anthropogenic marine debris  
Beach survey  
Volunteer participation  
Environmental education  
Plastics

### ABSTRACT

Anthropogenic marine debris (AMD) is an ubiquitous problem, which has motivated public participation in activities such as beach surveys and clean-up campaigns. While it is known that beaches in the SE Pacific are also affected by this problem, the quantities and types of AMD remain largely unknown. In the context of an outreach project, volunteers (~1500 high-school students) participated in a nation-wide survey of AMD on 43 beaches distributed randomly along the entire Chilean coast (18°S to 53°S). The mean density of AMD was 1.8 items m<sup>-2</sup> and the major types were plastics, cigarette butts and glass. Densities in central Chile were lower than in northern and southern Chile, which could be due to different attitudes of beach users or to intense beach cleaning in central regions. We suggest that public participation in surveys and cleaning activities will raise awareness and thereby contribute to an improvement of the situation.

© 2009 Elsevier Ltd. All rights reserved.

### 1. Introduction

Litter in the environment is a problem that affects beaches throughout the world (Derraik, 2002; Gregory and Andrady, 2003; Ivar do Sul and Costa, 2007). Anthropogenic marine debris (AMD) is associated with diverse human activities that might take place on the beach itself or far away in the case of floating marine debris, which is transported via marine currents towards the sea shore. The amounts of AMD have increased in a way that this is currently recognized as a worldwide problem (e.g. Frost and Cullen, 1997; Santos et al., 2008).

AMD causes particular concerns because its potential impacts on marine wildlife and biodiversity. Animals often confuse plastic bags, plastic pellets and Styrofoam with their food causing severe damage and even death (Sheavly, 2005; Mallory, 2008; Ryan, 2008). Floating marine debris may also serve as transport vector for a wide diversity of marine organisms, which rapidly colonize these substrata. Due to the high longevity of most plastic debris, these objects together with their inhabitants may travel over long distances along the sea surface, possibly invading new, previously uncolonized, habitats (e.g. Gregory, 1991; Winston et al., 1997). Besides the environmental impacts, AMD in the oceans can also cause problems to other human activities by fouling ship propellers

or clogging intake filters of powerplants or aquaculture systems (Sheavly, 2005). AMD can also negatively affect local tourism (Ballance et al., 2000; Sheavly, 2005).

In parallel to growing amounts of AMD, public awareness has also increased. This is reflected in beach surveys carried out in many countries of the world such as e.g. Jordan (Abu-Hilal and Al-Najjar, 2004), Japan (Kusui and Noda, 2003), Australia (Cunningham and Wilson, 2003) and Brazil (Oigman-Pszczol and Creed, 2007), among many others. Beach clean-ups involving many volunteer participants are commonly conducted on beaches throughout the world. This is the case of *The Ocean Conservancy*, an international organization that established the *Annual International Coastal Cleanup*, and *Clean Up the World*, an organization supported by the *United Nations Environment Programme* that also conducts frequent clean-up activities. Community participation in beach clean-up and survey activities has been valued by several authors who consider it as an important strategy of increasing public awareness (Rees and Pond, 1995; Jackson et al., 1997; Storrer and McGlashan, 2006). Volunteer participation also facilitates extensive samplings and thereby helps to obtain more information from a wider range of sites (Rees and Pond, 1995).

Even though AMD is a worldwide problem, it has been little studied in Latin America. In the SE Pacific region, there are only two countries (Colombia, Chile) for which studies about AMD in the environment have been published and these mainly focused on floating marine debris (Thiel et al., 2003; Ivar do Sul and Costa, 2007). Little is known about the reality of AMD on beaches from

\* Corresponding author. Address: Universidad Católica del Norte, Biología Marina, Larrondo 1281, Coquimbo, Chile. Tel.: +56 51 209939; fax: +56 51 209812.  
E-mail address: [thiel@ucn.cl](mailto:thiel@ucn.cl) (M. Thiel).

this part of the world and even less is known about the sources of this problem.


Beach surveys have been done in different parts of the world (e.g. Moore et al., 2001; Otley and Ingham, 2003; Shimizu et al., 2008). However, most of these studies are focused on limited sectors and often do not include many beaches. The present survey encompasses a range of 35° of latitude, covering a coastline >4000 km in the SE Pacific, where AMD was surveyed at 43 sites in a systematic and repeatable way. Knowledge of the abundances and types of AMD is important to identify possible sources, thereby facilitating the search for solutions (Santos et al., 2008). Also, being a cooperation between scientists and high-school students, such a public survey permits students, teachers and common people to visualize the dimensions of the AMD problem on their local beaches (Jackson et al., 1997). This raises their awareness about this problem, which helps to motivate them in getting involved with the possible solutions.

ches (Jackson et al., 1997). This raises their awareness about this problem, which helps to motivate them in getting involved with the possible solutions.

## 2. Materials and methods

### 2.1. Volunteer participation

We contacted social organizations, citizen groups and schools in cities and villages along the entire Chilean coast. At each institution that agreed to participate in the survey, one person acted as local coordinator of the activity. Sampling instructions and work sheets were sent to each local coordinator (information available at <http://www.cientificosdelabasura.cl/MuestreoNacionalguas.htm>).



**Fig. 1.** (A) Schematic overview of beach survey design. Station 1 (S1) corresponds to the “low station” and station 6 (S6) to the “high station” of each transect. (B) Example of a station being surveyed by volunteer participants. (C) The five geographic zones along the Chilean coast. Large roman numbers in the map refer to the administrative regions of Chile. Small Arabic numbers next to the map indicate survey sites.

Additionally, we provided printed guidelines with information about AMD and its impact on the marine environment. Booklets with an educational story informing about the fate of plastic bags in the marine environment were provided to motivate and inform participants; this booklet is directed at school kids from ages 5–15.

## 2.2. Beach surveys of anthropogenic debris

Beach surveys were conducted between August and September 2008 (i.e. during austral winter), and each local coordinator selected one or several beaches within the neighborhood of their school or organization. Most of the surveyed sites were sandy beaches, but some of them were pebble beaches; rocky shores were

not surveyed. At most sites, several transects were surveyed (up to 26 transects in Punta Arenas). These transects were perpendicular to the coastline, i.e. from the low tide line (low station) to the base of dunes (high station). On each transect a minimum of two stations were surveyed (between 2 and 6 stations, depending on the width of the beach) (Fig. 1a). Each station covered an area of  $3\text{ m} \times 3\text{ m}$ , which was delimited by ropes or measuring tape (Fig. 1b). All types of AMD within these  $9\text{ m}^2$  were counted and classified. This methodology is similar to the one that had been successfully used by Jackson et al. (1997) with students from the secondary school level (ages 15–17).

Data from each site were sent by the local coordinator via e-mail or regular mail to the coordinating office at Universidad

**Table 1**  
Institutions and numbers of volunteer participants in the survey of AMD in each administrative region from the Chilean coast. Names of local coordinators are also presented.

Zone	Region	City	School or institution	Number of students	Regional number of students	Coordinators
1	Arica y Parinacota Tarapacá	Arica	ONG Borde Río	23	23	José López
		Iquique	Colegio Maria Reina	71	71	Rossana Cáceres Norma Ponce
	Antofagasta	Tocopilla	Esc. Estados Unidos de Norteamérica	12	62	Sonia Arancibia
		Antofagasta	Esc. Diferencial Las Buganvillias	10		Julia Leo
2	Atacama	Chañaral	Colegio Chañares	40	103	–
		Chañaral	Esc. Diferencial José Luís Arancibia F103	15		Ernesto Tolmo Juan Carlos Pinto
	Coquimbo	Caldera	Escuela Villa las Playas	40	190	"Científicos de la Basura"
		Huasco	Escuela Byron Gigoux	25		
	Coquimbo	Coquimbo	Escuela Mireya Zuleta	23	190	"Científicos de la Basura"
		Coquimbo	Colegio Los Carrera	120		
	Coquimbo	Coquimbo	Colegio Santa María de Belén	120	190	"Científicos de la Basura"
		Coquimbo	Escuela La Herradura	120		
3	Valparaíso	La Serena	Colegio Los Héroes de la Concepción	35	76	Milenko Alvarado
		Los Vilos	Colegio Diego de Almagro	35		Nelson Vásquez
	Valparaíso	Puchuncaví	Escuela Básica de Maitencillo	40	76	Javier Travelli
		Zapallar	Escuela de Cachagua	36		Clara Köhnenkamp
	L. Bdo. O'Higgins	Navidad	Escuela Divina Gabriela	15	147	Ximena Morar
		Navidad	Escuela Fco. Chávez Cifuentes Rapel	17		
	L. Bdo. O'Higgins	Navidad	Liceo Pablo Neruda	28	147	Ximena Morar
		Navidad	Esc. Unión de Mujeres Americanas	17		
	L. Bdo. O'Higgins	Pichilemu	Charly's School	70	147	Vilma Leiva
		Pichilemu	Escuela Digna Camilo	70		
	Maule	Llico	Liceo Entre Aguas de Llico	24	73	Viviana Contreras
		Bolleruca	Escuela Ema Cornejo de Cardoén	20		
	Maule	Chanco	Escuela de Loanco	7	73	Viviana Contreras
		Chanco	Escuela de Loanco	7		
	Maule	Constitución	Escuela Barra Barrios Claudio Antonio	22	73	Viviana Contreras
		Concepción	Colegio Alemán de Concepción	140		
4	Bío-Bío	Concepción	Colegio Sagrados Corazones	38	263	Claudio Vergara
		Lebu	Escuela Fresia Graciela Muller	27		Jenny Soto
	Bío-Bío	Lebu	Escuela Fresia Graciela Muller	27	263	Mónica Rodríguez
		Tirúa	Liceo C90 Trapaqueante	14		Cynthia Valenzuela
	Bío-Bío	Tirúa	Liceo C90 Trapaqueante	14	263	Daniela Fuentealba
		Tirúa	Escuela Eloísa González F866	44		Orlando Cayupe
	La Araucanía	Carahue	Liceo Llaguey N° 51	40	65	Jorge Guzmán
		Carahue	Liceo Llaguey N° 51	40		Juan Francisco Gaete
	La Araucanía	Carahue	Liceo Llaguey N° 51	40	65	Juan Carlos Molina
		Carahue	Liceo Llaguey N° 51	40		Doris Salgado
	Los Ríos	Pto. Saavedra	Liceo Gabriela Mistral	25	61	Gabriela Quilodrán
		Valdivia	Escuela de Niebla	30		Martha Lidia Morales
	Los Ríos	Valdivia	Colegio Domus Mater	31	61	Silva Melo Necul
		Valdivia	Colegio Domus Mater	31		Olga San Martín
5	Los Lagos	Pto. Montt	Colegio de Lenguaje Amapola	50	63	Patricio Manríquez
		Quemchi	Escuela Borde Mar	13		Álvaro Volpi
	Aysén	Chacabuco	Escuela de Chacabuco	22	22	Alex Arismendi
		Chacabuco	Escuela de Chacabuco	22		Cristian Gaete
	Magallanes	Pto. Natales	Liceo Politécnico Luís Cruz Martínez	312	312	Gonzalo Hormazabal
		Pto. Natales	Liceo Politécnico Luís Cruz Martínez	312		Pablo Araya
	Magallanes	Pto. Natales	Colegio Puerto Natales	312	312	Pablo Araya
		Pto. Natales	Colegio Puerto Natales	312		
	Magallanes	Pto. Natales	Escuela Coronel Santiago Bueras	312	312	Pablo Araya
		Pto. Natales	Escuela Coronel Santiago Bueras	312		
	Magallanes	Pto. Natales	Escuela Capitán Juan Ladrillero	312	312	Pablo Araya
		Pto. Natales	Escuela Capitán Juan Ladrillero	312		
	Magallanes	Pto. Natales	Liceo Gabriela Mistral	312	312	Pablo Araya
		Pto. Natales	Liceo Gabriela Mistral	312		
	Magallanes	Pto. Natales	Colegio Maria Mazzarello	312	312	Pablo Araya
		Pto. Natales	Colegio Maria Mazzarello	312		
	Magallanes	Pto. Arenas	Escuela Bernardo O'Higgins	312	312	Pablo Araya
		Pto. Arenas	Escuela Bernardo O'Higgins	312		
	Magallanes	Pto. Arenas	Escuela España	312	312	Pablo Araya
		Pto. Arenas	Escuela España	312		
	Magallanes	Pto. Arenas	Grupo Scout Don Bosco	312	312	Pablo Araya
		Pto. Arenas	Grupo Scout Don Bosco	312		
	Magallanes	Pto. Arenas	Colegio Pierre Faure	312	312	Pablo Araya
		Pto. Arenas	Colegio Pierre Faure	312		
Total					1531	36

Católica del Norte in Coquimbo, where they were analyzed. In this study we only used data of AMD, excluding debris of natural origin. In order to explore regional variations in the densities of AMD the entire Chilean coast was subdivided into five zones (Fig. 1c). Zone 1 includes the extreme northern regions, which are characterized by a dry desert climate and the absence of large rivers. Zone 2 covers the northern-central region with a semi-arid climate, with some local rivers, while zone 3 corresponds to the central regions of the country with a Mediterranean climate. Zone 4 includes the southern region with a temperate-cold climate, high precipitation and large rivers, and zone 5 covers the extreme south regions, characterized by high rainfall, windy climate and deep fjords. The highest population density with intense tourism is found in zone 3, followed by zones 2 and 4. The beaches of zone 3 are in close reach of the metropolitan region of Santiago, where about 7 million people (half the population of Chile) are living.

### 2.3. Data evaluation

To determine whether densities of AMD differed significantly among the regions of the Chilean coast, we conducted a nonparametric Kruskal-Wallis analysis. When differences were detected, the nonparametric post-hoc Dunn's test for unequal sample size was applied. To examine whether the distribution of AMD within a beach differed, and whether this pattern was consistent along the Chilean coast, we compared the AMD densities of the highest and lowest station of each transect across the five zones. We applied a nested ANOVA, where stations (highest and lowest) were nested within the five zones. Since variances were not homogenous across the groups, data were transformed to the reciprocal root ( $1/\sqrt{x+1}$ ). This transformation method was selected after applying the protocol suggested by Fry (1993), which consists of several steps to select the most appropriate transformation method depending on the relationship between means and variances across the groups to be compared. After transformation, variances across the groups were found to be homogeneous (Bartlett's test,  $X^2 = 14.932$ ,  $p = 0.093$ ) and residuals were close to normality (Liliefors test  $p = 0.04$ ). To adjust for unequal sample size a General Linear Model (Zar, 1999) was applied using SYSTAT, which was used for all statistical analyses. When differences were detected (at  $\alpha = 0.05$ ), a post-hoc Tukey test was carried out to identify which zones and stations differed among each other. Finally, the means of AMD densities and standard deviations were back-transformed.

## 3. Results

### 3.1. Participation and characteristics of survey participants

A total of 1531 students from 46 schools covering the 14 coastal regions of Chile participated in the survey of AMD. The participating schools and groups were supported by 46 different institutions, including local city governments, corporations, private foundations and other NGO's. The highest number of participants was in the Magallanes region (312 volunteers), followed by the Bío-Bío (263) and Coquimbo regions (190) (Table 1).

For most volunteers this was the first time that they participated in a scientific survey. However, the local coordinators were familiar with the scientific method and in particular with quadrat counts (many of them had studied marine biology). When reporting their data, some of the survey participants also expressed surprise about the amounts of AMD found on their local beaches. In addition to the data, we had requested photographs of the beaches and the activity from the local coordinators. Without being particularly asked, many participants also inferred about the possible sources of AMD on their local beaches. For example, a group from northern

Chile associated the abundance of bottles and glass fragments with the carelessness of beach goers. Several local coordinators also mentioned that during the activity, participants started discussing about possible ways of mitigating the problem. These observations suggest that the sensitivity of the volunteers towards the problem of AMD on their local beaches increased during the participation in the survey.

### 3.2. Anthropogenic Marine Debris on the beaches in the SE Pacific

On all beaches surveyed in this study a total of 21,146 items of AMD were found and classified. The highest percentages of AMD on most beaches were plastics, cigarette butts and glass fragments. The proportions of the different types of AMD varied between the five geographic zones of the Chilean coast (Fig. 2). Plastics were very common in all five zones. In the northern zones (zones 1 and 2) cigarette butts predominated, while in zone 3 coal (classified under *other items*) was very common. In the two southern zones (4 and 5) metal, glass pieces, plastic and other items (mostly Styrofoam and cloths) abounded (Fig. 2).

The average density of AMD on beaches from the SE Pacific was  $1.8 \text{ items m}^{-2}$  (Fig. 3). Densities of AMD were significantly different between the regions along the Chilean coast (Kruskal-Wallis,  $H_c = 165.6$ ,  $p < 0.001$ ). Dunn's nonparametric multiple range test separated the regions into three statistically different groups: (i) Antofagasta; (ii) Tarapaca, Aysen and Magallanes and (iii) Arica and all other regions from Atacama to Los Lagos. The Antofagasta region featured the highest densities of AMD with most stations above  $2 \text{ items m}^{-2}$  and some stations even exceeding  $20 \text{ items m}^{-2}$





Fig. 2. Type and percentages of different categories of Anthropogenic Marine Debris in each of the five zones of the Chilean coast (average density of AMD for each zone is also presented). Numbers in parentheses correspond to the number of sites and stations surveyed in each zone. Black dots in the map indicate surveyed sites in each region.



**Fig. 3.** Density of AMD in each administrative region along the Chilean coast. The interrupted line indicates the global average of AMD density for the Chilean coast. Box plots show the mean, percentiles, range and outliers. Capital letters A–C indicate significant differences between regions (Dunn’s test,  $p < 0.05$ ). Numbers behind the name of each region correspond to the number of sites surveyed; the number of transects; number of stations.

$m^{-2}$ , clearly above the national average (Fig. 3). In the group of regions with low AMD densities, the regions of Atacama and Araucanía were the ones with the lowest densities, not exceeding 2

**Table 2**

Results of nested ANOVA for mean densities of AMD transformed to the square root between zones and among stations (highest and lowest) within a zone.


Source	Type III SS	df	MS	F-ratio	p-value
Zone	3.225	4	0.806	16.198	<0.001
Station (zone)	1.900	5	0.380	7.632	<0.001
Error	24.592	494	0.050	–	–

items  $m^{-2}$  at most stations. However, in some of the regions within this group, there were also stations with AMD densities above 10 items  $m^{-2}$ . Of the 43 surveyed sites along the country, the cleanest beaches of Chile were Playa Grande from the Atacama region in zone 2, Playa Tirúa and Playa Cura from the Bío-Bío region in zone 4, with an average of 0.3 items  $m^{-2}$ . Despite low densities, also on those beaches plastics are the most common AMD.

Mean AMD densities differed between zones and also among stations within a zone (Table 2). The post-hoc Tukey test indicated that the mean of zone 1 was significantly higher than that of the other zones (Fig. 4). The lowest abundances were reported from zones 2 and 3. Significant differences were found between the high and low stations in zones 1, 2 and 5, but not between the high and low stations in zones 3 and 4 (Tukey’s test; Fig. 4). At most of the studied sites the highest densities of AMD were found at the upper transect stations with an overall average of 2.1 items  $m^{-2}$ . This average was lower at the lowest station of each transect, i.e. close to the water line (1.7 items  $m^{-2}$ ).

**4. Discussion**

The national survey of AMD on beaches from the SE Pacific was well received by the volunteer participants, indicating that many people are concerned about the problem and willing to do something about it. All participants recognized and valued that this activity was not a clean-up campaign but rather a survey to provide a first diagnosis. The participation of experienced local coordinators ensured that data were comparable across the surveyed sites and zones, despite the different backgrounds of the volunteer participants.



**Fig. 4.** Density of AMD (Mean + Standard deviation, back-transformed) at the low and high stations in the five zones along the Chilean coast. Capital letters A–C indicate significant differences between zones, stars (\*) indicate significant differences between high and low stations within a zone (Tukey,  $p < 0.05$ ).  $n$  = Number of transects surveyed in each zone. Small insert shows a schematic cross section of a beach transect showing the low and high station (numbers of intermediate stations was variable at the different sites, which is why we only considered the highest and the lowest station for this analysis).

The analysis of the data indicated a relatively uniform distribution of AMD across the beaches from the SE Pacific with some regional variations, the reasons for which will be discussed in the following.

#### 4.1. Volunteer participation in AMD survey – data reliability

The present study employed a similar methodology as previously used in other surveys of AMD on local beaches (e.g. Jackson et al., 1997). As highlighted by Rees and Pond (1995), this kind of study has the potential to ensure wide geographic coverage without the necessity of extensive financial support, because it is based on the help of volunteer participants. Our study confirmed this, because with the aid of volunteers a large part of the SE Pacific could be covered. A synchronized survey, such as the one reported herein, would have been virtually impossible without the help of volunteers.

Materials used for this survey were simple (lines to delimit the 9 m<sup>2</sup> area at each station, gloves for sorting the AMD and paper forms), and most of these items were available in the local schools or could be acquired at moderate costs. In some cases transporting volunteers to the beaches and mailing the filled forms were among the major costs, which usually were assumed by the schools. The minimal requirements for equipment and transport have been previously emphasized as a major advantage of volunteer-based beach surveys (Rees and Pond, 1995).

In general, since most items that make up the bulk of AMD are very common in any household garbage bin, it is relatively easy even for untrained volunteers to classify and count the principal types of AMD. However, with the purpose of improving data quality, volunteers were also provided with a chart that highlighted the different types of AMD. Detailed and easily understandable instructions and guidance are considered essential in volunteer-based surveys (Rees and Pond, 1995). These instructions helped to avoid classification errors that could foul the collected data. Herein we only focused on AMD and we excluded natural marine debris such as e.g. wood, because these could either be of anthropogenic (manufactured) or natural (branches, trees) origin, thereby potentially introducing confusion (Alkalay et al., 2007).

One of the main concerns of surveys conducted with the help of volunteer participants is whether the data are reliable and comparable. For example, Moore et al. (2001) conducted a carefully designed study on litter densities from beaches in southern California, and they then compared their data with those obtained from volunteers during a beach clean-up campaign. Densities calculated from volunteer data were 50-times lower than densities based on their own data, which they attributed to the fact that: (i) volunteers focused on cleaning up the beach rather than on sur-

veying densities and (ii) collected mainly larger items (Moore et al., 2001). In the instructions to our local coordinators, we emphasized that the present study was not a clean-up activity, but rather a survey of litter densities. This message was repeated in both verbal and written communications prior to the survey, and the photographic and written documentation that we received together with the data confirmed that all participants carefully adhered to these instructions. Thus, we consider that the data obtained in this study reliably reflect the densities of AMD on the surveyed beaches.

One potential problem of public beach surveys is the bias at the time of selecting a beach for the activity. This decision could be biased towards easy access or more contaminated beaches (Rees and Pond, 1995). Herein, the local coordinators usually chose the beaches according to beach access (i.e. walking distance to the school or availability of public transport), and preconceptions about the amounts of AMD were not part of the decision process. Overall, the nation-wide survey includes all types of beaches (i.e. within and outside of large cities, contaminated and relatively clean), thereby ensuring the representativeness of the data.

#### 4.2. Beach debris in Chile and its origins

Following the global tendency (Derraik, 2002), plastics were among the most common debris types on almost all surveyed beaches in Chile. Plastics are used in almost all human activities (professional and recreational), and they are a very resistant material, persisting for long times in the environment, both under water as well as in the air (Pegram and Andrady, 1989). In general, in many parts of the world, and in particular on beaches used by tourists, a high proportion of AMD comes from beach users themselves (e.g. Abu-Hilal and Al-Najjar, 2004; Ariza et al., 2008). Several indicators, such as the specific composition of AMD and the distribution pattern on the beach, support this suggestion. For example, in some of the surveyed beaches glass fragments (mainly bottles) and cigarette butts were very common, which are two types of AMD usually directly attributed to local beach users (Silva-Iñiguez and Fischer, 2003; Taffs and Cullen, 2005). Similar as in the northern and central zones, Silva-Iñiguez and Fischer (2003) observed highest densities of AMD in the upper zones of the beach, a situation also reported from other parts of the world (Jackson et al., 1997; Cunningham and Wilson, 2003; Claereboudt, 2004). This distribution pattern within a beach suggests local sources of AMD, because beach goers typically frequent the highest parts of the beach. Additionally, AMD spread over the beach might be reshuffled by tides and waves, concentrating it high up on the beach (Taffs and Cullen, 2005).

**Table 3**

Densities of AMD reported from beaches throughout the world.

Country	Number of surveyed beaches	Average densities (items m <sup>-2</sup> )	Maximum densities (items m <sup>-2</sup> )	Reference
Scotland	16	0.4	2.3	Velander and Mocogni (1999)
Brazil	2	0.7	2.1	Araújo et al. (2006)
Brazil	10	0.14	~0.5	Oigman-Pszczol and Creed (2007)
Japan <sup>a</sup>	34	45,000	280,000	Fujieda and Sasaki (2005)
Japan <sup>a</sup>	18	3.4	2200	Kusui and Noda (2003)
Russia	8	0.2	16.7	Kusui and Noda (2003)
Oman	11	~0.4	~0.9	Claereboudt (2004)
Jordan	3	4.0	7.4	Abu-Hilal and Al-Najjar (2004)
Panama	19	3.6	–	Garrity and Levings (1993)
Australia	1	0.5	0.5	Foster-Smith et al., 2007
Australia	6	0.1	0.3	Cunningham and Wilson (2003)
Israel	6	–	0.9	Bowman et al. (1998)
Pitcairn Islands	2	0.2	0.4	Benton (1995)
Ireland	1	0.2	–	Benton (1995)
Indonesia	21	4.6	–	Evans et al. (1995)
Chile	43	1.8	82.7	This study

<sup>a</sup> These studies counted individual pellets of fragmented Styrofoam, an item usually not counted in most other studies.



of AMD on local beaches, as also suggested by Araújo and Costa (2007) and Santos et al. (2008) for some beaches in Brazil. The input of AMD by rivers could be intensified by rain (Shimizu et al., 2008), and since the south of Chile is a rainy region, river input might be locally significant (see very high AMD density in one station of the region Los Ríos in Fig. 3). In the fjord region of southern Chile (zone 5), aquaculture activities appear to be another important source of AMD on local beaches (Hinojosa and Thiel, 2009).

Densities of AMD on beaches in the SE Pacific are similar to those reported from other parts of the world (Table 3). Even though the population density along the Chilean coast is much lower than in many other regions of the world, the densities of AMD on local beaches are comparable to those found in densely populated areas of the northern hemisphere. Since there exists strong indication that most AMD on beaches from the SE Pacific has local sources, the solutions should also be sought at the local level.

#### 4.3. Volunteer participation

Most volunteers participating in this survey were students from the middle and high-school level (ages 10–16). The survey was part of an outreach project introducing the scientific method to school kids. While first contacts were made with the teachers and these took the decision to participate, students generally participated enthusiastically in the survey (Fig. 5a and b). One major feature of this survey is that students did not only repeat simple scientific experiments but they produced real data, which would not have been available otherwise. They were aware of this throughout and they also knew that they were part of a larger survey. At present, we do not know to which degree this knowledge might have fostered their willingness to participate, but the positive responses we received from the participating schools suggest that this could be a very promising model. Collaborative studies between scientists and school students such as this survey of AMD could be a powerful tool in increasing sample size and geographic coverage (see also Rees and Pond, 1995). There is a wealth of questions from all branches of natural sciences (biology, environmental ecology, geography, sociology) that could be tackled by scientist–teacher–student teams.

The call for volunteers in the beach survey was positively received by the contacted organizations, institutions and schools. Most people immediately declared their willingness to participate in this activity. This shows the potential of involving volunteers in activities related to environmental protection, which had also been highlighted by other authors (Jackson et al., 1997; The Ocean Conservancy, 2006; Storrer and McGlashan, 2006). Herein, volunteers were introduced to a common environmental problem, and by actively participating in the survey and noticing the large amounts of AMD, their awareness of the problem increased. The involvement of the general public in this kind of projects is considered very important at the moment of teaching citizens about the problem and impacts of AMD (Earll et al., 2000).

The volunteers participating in this study recognized, in an autonomous way, the causes that provoke the debris problem. This kind of discussion proves that volunteers became more sensitive to the problem of AMD at the end of the activity, an observation also highlighted by other authors (Jackson et al., 1997; Storrer and McGlashan, 2006). Publicity is considered an important part in the search for solutions (Velandar and Mocogni, 1998). In this study, the preliminary results of the survey were published in national and local newspapers (Fig. 5c), thereby also reaching many people who did not actively participate in the survey. Future beach surveys and clean-up activities will show whether this type of publicity indeed helped raise awareness.

#### 4.4. Conclusions and suggestions

This study is only the first step in revealing the reality of AMD on beaches of the SE Pacific. The obtained results can, however, already be used to mitigate the AMD problem in the SE Pacific. The causes for the high densities of AMD on the surveyed beaches are variable and often depend on local realities, including: (i) lack of conscience by beach users, (ii) insufficient garbage bins, (iii) no recycling and reutilization system, (iv) absence of inspection, (v) lack of information about the extent of the problem. Many of these causes are transversal and valid for anthropogenic debris in any environment. The principal solutions are better infrastructure and environmental education. Starting in local schools and getting the general public involved in these activities (Fig. 5d) appear to be some of the most promising avenues in resolving the common problem of AMD.

#### Acknowledgements

We are grateful to all volunteers along the country and in particular we thank the local coordinators. This survey would not have been possible without their help, and hopefully this publication helps that all of us can enjoy cleaner beaches in the future. This project was funded by EXPLORA-CONICYT (ED12070) and Universidad Católica del Norte, Chile. Further information and more details can be found at: [www.cientificosdelabasura.cl](http://www.cientificosdelabasura.cl).

#### References

- Abu-Hilal, A.H., Al-Najjar, T., 2004. Litter pollution on the Jordanian shores of the Gulf of Aqaba (Red Sea). *Marine Environmental Research* 58, 39–63.
- Alkalay, R., Pasternak, G., Zask, A., 2007. Clean-coast index – a new approach for beach cleanliness assessment. *Ocean and Coastal Management* 50, 352–362.
- Araújo, M.C.B., Costa, M.F., 2007. Visual diagnosis of solid waste contamination of a tourist beach: Pernambuco, Brazil. *Waste Management* 27, 833–839.
- Araújo, M.C.B., Santos, P.J.P., Costa, M.F., 2006. Ideal width of transects for monitoring source-related categories of plastics on beaches. *Marine Pollution Bulletin* 52, 957–961.
- Ariza, E., Jiménez, J.A., Sardá, R., 2008. Seasonal evolution of beach waste and litter during the bathing season on the Catalan coast. *Waste Management* 28, 2604–2613.
- Ballance, A., Ryan, P.G., Turpie, J.K., 2000. How much is a clean beach worth? The impact of litter on beach users in the Cape Peninsula, South Africa. *South African Journal of Science* 96, 210–213.
- Barragán, J.M., Castro, C., Alvarado, C., 2005. Towards integrated coastal zone management in Chile. *Coastal Management* 33, 1–24.
- Benton, T.G., 1995. From castaways to throwaways: marine litter in the Pitcairn Islands. *Biological Journal of the Linnean Society* 56, 415–422.
- Bowman, D., Manor-Samsonov, N., Golik, A., 1998. Dynamics of litter pollution of Israeli Mediterranean beaches: a budgetary, litter flux approach. *Journal of Coastal Research* 14, 418–432.
- Claereboudt, M.R., 2004. Shore litter along sandy beaches of the Gulf of Oman. *Marine Pollution Bulletin* 49, 770–777.
- Cunningham, D.J., Wilson, S.P., 2003. Marine debris on beaches of the greater Sydney Region. *Journal of Coastal Research* 19, 421–430.
- Derraik, J.G.B., 2002. The pollution of the marine environment by plastic debris: a review. *Marine Pollution Bulletin* 44, 842–852.
- Earll, R.C., Williams, A.T., Simmons, S.L., Tudor, D.T., 2000. Aquatic litter, management and prevention – the role of measurement. *Journal of Coastal Conservation* 6, 67–78.
- Evans, S.M., Dawson, M., Day, J., Frid, C.L.J., Gill, M.E., Pattisina, L.A., Porter, J., 1995. Domestic waste and TBT pollution in coastal areas of Ambon Island (Eastern Indonesia). *Marine Pollution Bulletin* 30, 109–115.
- Foster-Smith, J., Birchenough, A.C., Evans, S.M., Prince, J., 2007. Human impacts on Cable Beach, Broome (Western Australia). *Coastal Management* 35, 181–194.
- Frost, A., Cullen, M., 1997. Marine debris on northern New South Wales beaches (Australia): sources and the role of beach usage. *Marine Pollution Bulletin* 34, 348–352.
- Fry, J.C., 1993. One-way analysis of variance. In: Fry, J.C. (Ed.), *Biological Data Analysis. A Practical Approach*. Oxford University Press, pp. 3–39.
- Fujieda, S., Sasaki, K., 2005. Stranded debris of foamed plastic on the coast of ETA Island and Kurahashi Island in Hiroshima Bay. *Nippon Suisan Gakkaishi* 71, 755–761.
- Garrity, S.D., Levings, S.C., 1993. Marine debris along the Caribbean coast of Panama. *Marine Pollution Bulletin* 26, 317–324.

- Gregory, M.R., 1991. The hazards of persistent marine pollution: drift plastics and conservation islands. *Journal of the Royal Society of New Zealand* 21, 83–100.
- Gregory, M.R., Andrady, A.L., 2003. Plastics in the marine environment. In: Andrady, A.L. (Ed.), *Plastics and the Environment*. John Wiley and Sons, Inc., pp. 379–401.
- Hinojosa, I.A., Thiel, M., 2009. Floating marine debris in fjords, gulfs and channels of southern Chile. *Marine Pollution Bulletin* 58, 341–350.
- Ivar do Sul, J.A., Costa, M.F., 2007. Marine debris review for Latin America and the Wider Caribbean Region: from the 1970s until now, and where do we go from here? *Marine Pollution Bulletin* 54, 1087–1104.
- Jackson, N.L., Cerrato, M.L., Elliot, N., 1997. Geography and fieldwork at the secondary school level: an investigation of anthropogenic litter on an estuarine shoreline. *Journal of Geography* 96, 301–306.
- Kusui, T., Noda, M., 2003. International survey on the distribution of stranded and buried litter on beaches along the Sea of Japan. *Marine Pollution Bulletin* 47, 175–179.
- Mallory, M.L., 2008. Marine plastic debris in northern fulmars from the Canadian high Arctic. *Marine Pollution Bulletin* 56, 1486–1512.
- Martínez, E., Maamaatuaiahutapu, K., Taillandier, V., 2009. Floating marine debris surface drift: convergence and accumulation toward the South Pacific subtropical gyre. *Marine Pollution Bulletin*, doi:10.1016/j.marpolbul.2009.04.022.
- Moore, S.L., Gregorio, D., Carreon, M., Weisberg, S.B., Leecaster, M.K., 2001. Composition and distribution of beach debris in Orange County, California. *Marine Pollution Bulletin* 42, 241–245.
- Oigman-Pszczol, S.S., Creed, J.C., 2007. Quantification and classification of marine litter on beaches along Armacao dos Búzios, Rio de Janeiro, Brazil. *Journal of Coastal Research* 23, 421–428.
- Otley, H., Ingham, R., 2003. Marine debris surveys at Volunteer Beach, Falkland Islands, during the summer of 2001/02. *Marine Pollution Bulletin* 46, 1534–1539.
- Pegram, J.E., Andrady, A.L., 1989. Outdoor weathering of selected polymeric materials under marine exposure conditions. *Polymer Degradation and Stability* 26, 333–345.
- Rees, G., Pond, K., 1995. Marine litter monitoring programmes – a review of methods with special reference to national surveys. *Marine Pollution Bulletin* 30, 103–108.
- Ryan, P.G., 2008. Seabirds indicate changes in the composition of plastic litter in the Atlantic and south-western Indian Oceans. *Marine Pollution Bulletin* 56, 1406–1409.
- Santos, I.R., Friedrich, A.C., Wallner-Kersanach, M., Fillman, G., 2005. Influence of socio-economic characteristics of beach users on litter generation. *Ocean and Coastal Management* 48, 742–752.
- Santos, I.R., Friedrich, A.C., Ivar do Sul, J.A., 2008. Marine debris contamination along undeveloped tropical beaches from northeast Brazil. *Environmental Monitoring and Assessment* 148, 455–462.
- Sheavly, S.B., 2005. Sixth Meeting of the UN Open-ended Informal Consultative Process on Oceans and the Law of the Sea. *Marine Debris – An Overview of a Critical Issue for Our Oceans*. 6–10 June 2005. <[http://www.un.org/Depts/los/consultative\\_process/consultative\\_process.htm](http://www.un.org/Depts/los/consultative_process/consultative_process.htm)>.
- Shimizu, T., Nakai, J., Nakajima, K., Kozai, N., Takahashi, G., Matsumoto, M., Kikui, J., 2008. Seasonal variations in coastal debris on Awaji Island, Japan. *Marine Pollution Bulletin* 57, 182–186.
- Silva-Iñiguez, L., Fischer, D.W., 2003. Quantification and classification of marine litter on the municipal beach of Ensenada, Baja California, Mexico. *Marine Pollution Bulletin* 46, 132–138.
- Somerville, S.E., Miller, K.L., Mair, J.M., 2003. Assessment of the aesthetic quality of a selection of beaches in the Firth of Forth, Scotland. *Marine Pollution Bulletin* 46, 1184–1190.
- Storrier, K.L., McGlashan, D.J., 2006. Development and management of a coastal litter campaign: the voluntary coastal partnership approach. *Marine Policy* 30, 189–196.
- Taffs, K.H., Cullen, M.C., 2005. The distribution and abundance of beach debris on isolated beaches of northern New South Wales, Australia. *Australasian Journal of Environmental Management* 12, 244–250.
- The Ocean Conservancy, 2006. *A World of Difference*. The International Coastal Cleanup 2006, p. 39.
- Thiel, M., Hinojosa, I., Vásquez, N., Macaya, E., 2003. Floating marine debris in coastal waters of the SE-Pacific (Chile). *Marine Pollution Bulletin* 46, 224–231.
- Uneputty, P.A., Evans, S.M., 1997. Accumulation of beach litter on Islands of the Pulau Seribu Archipelago, Indonesia. *Marine Pollution Bulletin* 34, 652–655.
- Velander, K., Mocogni, M., 1998. Maritime litter and sewage contamination at Cramond Beach Edinburgh – a comparative study. *Marine Pollution Bulletin* 36, 385–389.
- Velander, K.A., Mocogni, M., 1999. Beach litter sampling strategies: is there a 'best' method? *Marine Pollution Bulletin* 38, 1134–1140.
- Winston, J.E., Gregory, M.R., Stevens, L.M., 1997. Epibionts, and other biota associated with pelagic plastics. Encrusters a review of biogeographical, environmental, and conservation issues. In: Coe, J.M., Rogers, D.B. (Eds.), *Marine debris: sources, impacts, and solutions*. Springer-Verlag, New York, USA, pp. 81–97.
- Zar, J.H., 1999. *Biostatistical Analysis*. Prentice Hall, New Jersey, USA. 663p.